








Large Scale Structure Simulations


Katrin Heitmann

Argonne National Laboratory High Energy Physics/ Mathematics & Computer Science Division





The Role of N-body Simulations in Cosmology




All detailed knowledge of structure formation in the nonlinear regime comes from N-body simulations

- Discovering new physics in the nonlinear regime
- Exploring new cosmological probes
- Analysis of survey results with precision predictions
- Solving cosmological inverse problems
- Basis for generating mock observations
- Estimation of measurement errors



The Role of N-body Simulations in Cosmology



All detailed knowledge of structure formation in the nonlinear regime comes from N-body simulations

- Discovering new physics in the nonlinear regime
- Exploring new cosmological probes
- Analysis of survey results with precision predictions
- Solving cosmological inverse problems
- Basis for generating mock observations
- Estimation of measurement errors

Risa's talk



N-body Simulation Basics


- Follow evolution of dark matter tracer particles in an expanding universe
- Solve Vlasov-Poisson equation
- In principle: N² problem (N = number of particles), in practice, approximate methods based on particle and grid solvers
 - Tree methods: Gadget, HOT, PKDGRAV
 - Adaptive mesh refinement: ART, Enzo, FLASH, Nyx, Ramses
 - Hybrid: Gadget-2, TreePM, HACC

$$\frac{\partial f_i}{\partial t} + \dot{\mathbf{x}} \frac{\partial f_i}{\partial \mathbf{x}} - \nabla \phi \frac{\partial f_i}{\partial \mathbf{p}} = 0, \quad \mathbf{p} = a^2 \dot{\mathbf{x}},$$

$$\nabla^2 \phi = 4\pi G a^2 (\rho(\mathbf{x}, t) - \langle \rho_{\rm dm}(t) \rangle) = 4\pi G a^2 \Omega_{\rm dm} \delta_{\rm dm} \rho_{\rm cr},$$

$$\delta_{\rm dm}(\mathbf{x}, t) = (\rho_{\rm dm} - \langle \rho_{\rm dm} \rangle) / \langle \rho_{\rm dm} \rangle),$$

$$\rho_{\rm dm}(\mathbf{x}, t) = a^{-3} \sum_{i} m_i \int d^3 \mathbf{p} f_i(\mathbf{x}, \dot{\mathbf{x}}, t).$$





The Virtues of N-body Simulations

Advantages:


- Capture relevant physics well out into the nonlinear regime
- Cover large dynamic range: from Gigaparsecs down to less than a kiloparsec, therefore allows coverage of survey size volumes
- Underlying physics is well understood
- Several proven numerical paradigms exist, these allow exploration of new highperformance architectures
- No free physics parameters, code parameters can be set to reach sub-percent accuracy
- Simulations can be carried out relatively fast (days to weeks time scale), therefore allow exploration of cosmological parameter space, many realizations can be carried out
- Post-processing via halo occupation distribution models or semi-analytic models allows exploration of additional physics in a straightforward way

Disadvantage:

Miss physics on small scales (gas dynamics, feedback..., see Nick's talk),
 require phenomenological techniques based on observations (self-calibration)




Physics Results on All Scales







Surveys Need Statistics!













Discovering New Physics in the Nonlinear Regime


- Halo profiles: Navarro, Frenk & White (NFW, 1996, 1997) find that the density profile of a halo as a function of radius has the same shape independent of mass and cosmology
- Universality of the mass function: For a particular halo definition (friends-of-friends with a linking length of b=0.2) and expressed via the variance of the linear density field σ²(z,M), mass function is independent of epoch/cosmology at the 20% level
- Non-gaussianity/bias: Non-gaussianity effects clustering of dark matter halos, bias is strongly scale dependent








Exploring New (and Established) Cosmological Probes

- Sensitivity of cosmological probes to new physics (e.g. early dark energy, modified gravity)
- Estimate of the strength of expected signal over noise
- New cosmological probes













Analysis of Survey Results: Three Examples

- Cluster Cosmology: Cosmological constraints from Chandra X-ray cluster observations, mass function matched to Tinker et al. 2008 fit from N-body simulations
- Weak lensing: Cosmological constraints from Canada-France-Hawaii-Telescope Legacy Survey (CFHTLS), Markov Chain Monte Carlo calculations based on Smith (Halo) fit for nonlinear power spectrum (calibrated with N-body simulations)
- Small scale galaxy clustering: Cosmological constraints from Sloan Digital Sky Survey, modeling based on Halo Occupancy Distribution (HOD), includes calibration of mass function, power spectrum, bias from N-body simulations




Solving the Inverse Problem

- Challenge: To extract cosmological constraints from observations in nonlinear regime, need to run Marko Chain Monte Carlo code; input: 10,000 -100,000 different models
- Current strategy: Fitting functions for e.g. P(k), accurate at 10% level, not good enough!
- Brute force: Simulations, ~30 years on 2000 processor cluster...
- Only alternative: emulators



Heitmann et al. 2006, Habib et al. 2007




Cosmic Emulator in Action: LSSFast


- Instantaneous 'oracle' for nonlinear power spectrum, reduces compute time from weeks to negligible, accurate at 1% out to k~1/Mpc for wCDM cosmologies
- Enables direct MCMC with results from full simulations for the first time



Hybrid Simulation Codes: HACC and TreePM

- Hardware Accelerated Cosmology Code (HACC) (Habib et al. 2009, Pope et al. 2010)
 - New cosmology code designed with major aim of easy portability between different high-performance computing architectures
 - Long-range force: particle mesh, short-range force: particle-particle or tree, architecture dependent
 - Strong collaboration between Los Alamos and Argonne National Labs in code development
 - First version ran on Roadrunner, Cell-accelerated machine, second version on GPU cluster (OpenCL implementation); at SC10: run on heterogenous system with in-situ visualization, now: port to BlueGene system
 - First science: BAO in the Lyman-alpha forest (9 PM runs, 64 billion particles each), precision cosmology with high-resolution code (16 billion particles and up)
- **TreePM** (White 2002)
 - Well established N-body code, used for very large number of science projects: weak lensing, galaxy clustering, BAO, mock catalogs, ...






Halos from Roadrunner simulation, visualization: J. Woodring



Adaptive Mesh Simulation Codes


- Nyx: New adaptive mesh refinement (AMR) code developed at LBNL
- Block-structured AMR with multigrid Poisson solver
- Main focus for AMR codes (also Enzo, ART): hydrodynamics simulations, discussed in detail in Nick's talk





Analysis Capabilities: ParaView and yt

- Analysis of simulations as demanding as carrying out the simulations
- Visualization-aided analysis
 - Visualization is important for understanding the results, and for verification & validation
 - Visualization and analysis have to go hand-in-hand to ensure efficiency
- ParaView (open source)
 - Based on VTK (Visualization Tool Kit), cosmological analysis capabilities focusing on particle data have been developed (LANL/ANL collaboration, J. Ahrens, J. Woodring, P. Fasel, K. Heitmann, S. Habib, J. Insley...)
 - Parallel reader, halo finder, sub-sampler, embedded plotting and analysis routines
- yt (open source)
 - Focus on astrophysical data, AMR, particles also supported, started by Enzo community (SLAC), expanding fast
 - Parallel halo finder, merger trees, ...
 - Lives also within ParaView now







Some Recent Science Highlights

- Collaboration members have very successful track record in using N-body simulations to explore new physics and analyze survey data
- Cross-laboratory collaborations are established, more being planned
- World-leading capabilities exist

Galaxy Surveys

- The clustering of massive galaxies at z~0.5 from the first semester of BOSS data, M.
 White et al. ApJ 728, 126 (2011), LBNL
- Cosmological constraints from galaxy clustering and the the mass-to-number ratio of galaxy clusters, J. Tinker, E. Sheldon, R. Wechsler et al., ApJ (2011), BNL/SLAC
- Statistics of satellite galaxies around Milky Way-like hosts, M. Busha, R. Wechsler et al, ApJ (2011), SLAC
- The real-space clustering of luminous red galaxies around z<0.6 quasars in the Sloan Digital Sky Survey, N. Padmanabhan, M. White, et al., MNRAS 397, 1862 (2009), LBNL

Weak Lensing

- Probing cosmology with weak lensing peak counts, J. Kratochvil, Z. Haiman, M. May, Phys. Rev. D81, 043519 (2010), BNL
- Cosmological information in weak lensing peaks, X. Yang, J. Kratochvil, S. Wang, E. A. Lim, Z. Haiman, M. May, Phys. Rev. D84, 043529 (2011), BNL

Some Recent Science Highlights

Baryon Acoustic Oscillations

- Constraining anisotropic baryon oscillations, N. Padmanabhan, M. White, Phys. Rev. D77, 123540 (2008), LBNL
- Non-linear structure formation and the acoustic scale, H.-J.Seo, E.Siegal, D.Eisenstein, M.White, ApJ 686, 13 (2008), LBNL
- Calibrating the baryon oscillation ruler for matter and halos, N. Padmanabhan, M. White, Phys. Rev. D80, 063508 (2009), LBNL
- Baryon Acoustic Oscillations in the Lyman alpha Forest
 - The acoustic peak in the Lyman alpha forest, A. Slosar, S. Ho, M. White, T. Louis, JCAP 10, 019 (2009), BNL/LBNL
 - Particle mesh simulations of the Lyman-alpha forest and the signature of baryon acoustic oscillations in the intergalactic medium, M. White, A. Pope, J. Carlson, K. Heitmann, S. Habib, P. Fasel, D. Daniel, Z. Lukic, ApJ 713, 383 (2010), LBNL/ANL

Redshift Space

- Testing cosmological structure formation using redshift-space distortions, W. Percival, M. White, MNRAS 393, 297 (2009), LBNL
- Towards an accurate model of the redshift space clustering of halos in the quasilinear regime, B. Reid, M. White, MNRAS, in press, LBNL



Some Recent Science Highlights

Clusters and Groups

- Mass function predictions beyond LCDM, S. Bhattacharya, K. Heitmann, M. White, Z. Lukic, C. Wagner, S. Habib, ApJ 732, 122 (2011), ANL/LBNL
- Cosmological constraints from the SDSS maxBCG cluster catalog, E. Rozo, R. Wechsler et al. ApJ, 708, 645 (2010), SLAC
- The structure of halos: Implications for group and cluster cosmology, Z. Lukic, D. Reed, S. Habib, K. Heitmann, ApJ 692, 217 (2009), ANL


Theory


- The Coyote Universe III: Simulation suite and precision emulator for the nonlinear matter power spectrum, E. Lawrence, K. Heitmann, M. White, D. Higdon, C. Wagner, S. Habib, B. Williams, ApJ 713, 1322 (2010), ANL/LBNL
- The Coyote Universe I: Precision determination of the nonlinear matter power spectrum, K. Heitmann, M. White, C. Wagner, S. Habib, D.Higdon, ApJ 713, 1322 (2010), ANL/LBNL
- The Cosmic Code Coparison Project, K. Heitmann, Z. Lukic, P. Fasel, S. Habib, M. Warren, M. White, J. Ahrens, L. Ankeny, R. Armstrong, B. O'Shea, P. Ricker, V. Springel, J. Stadel, H. Trac, Compt. Sci. Dis. 1, 015003 (2008), ANL/LBNL




Challenges for the Future

- Large cosmological volumes with ~ trillion particles are needed for ongoing and future large scale surveys
- Hundreds to thousands of realizations, not only for one cosmology!
- Coverage of cosmological model space
- Error control in simulations
- Supercomputer architectures are changing!
 Codes need to be flexible, close
 collaboration with computer science
 community very beneficial
- Simulation analysis, close collaboration with CS community very beneficial
- Data serving, close collaboration with CS community very beneficial
- Outstanding opportunity for HEP and ASCR to work close together and make an impact!
- National Laboratories poised to carry out an effort that can benefit the community









- N-body simulations hold a wealth of information about large-scale structure formation and probes of the dark Universe
- Precision predictions possible, important for analysis of large scale structure surveys
- Exciting opportunities for next-generation high-performance computing architectures (we are getting exa-scale ready!)